

Herpetofauna, municipality of Porto Murtinho, Chaco region, state of Mato Grosso do Sul, Brazil

Franco L. Souza^{1*}, Masao Uetanabaro², Paulo Landgref-Filho³, Liliana Piatti¹ and Cynthia P. A. Prado⁴

1 Universidade Federal de Mato Grosso do Sul, Centro de Ciências Biológicas e da Saúde, Departamento de Biologia. CEP 79070-900. Campo Grande, MS, Brazil.

2 Rua Clóvis, 24. CEP 79022-071. Campo Grande, MS, Brazil.

3 Rua João Cadete, 55. CEP 79044-370. Campo Grande, MS, Brazil.

4 Universidade Estadual Paulista, Departamento de Morfologia e Fisiologia Animal, Faculdade de Ciências Agrárias e Veterinárias. CEP 14884-900. Jaboticabal, SP, Brazil.

* Corresponding author. E-mail: flsouza@nin.ufms.br

ABSTRACT: Among the physiographic regions included in the central South American diagonal of open formations, the Chaco, with some endemic species, represents the southernmost dry area. In Brazil, the Chaco is found in southwestern corner of Mato Grosso do Sul state, mostly in the municipality of Porto Murtinho along the Paraguay and Apa rivers. From February 2008 to December 2009, we carried out an inventory of amphibians and reptiles in Porto Murtinho, using pitfall traps, time-limited searches, and occasional encounters. A total of 34 amphibian and 39 reptile species were registered. Although some typical Chacoan species were found, most of the species are open area dwellers that also occur in other open biomes, such as the Cerrado and Caatinga.

INTRODUCTION

Physiographic regions in the state of Mato Grosso do Sul consist of an array of open habitats, such as the Pantanal, Cerrado, Chaco, and dry forests with a distinct biota, formed by an interconnected mosaic of habitats. However, during the last few decades, most of this heterogeneous landscape has been converted into agricultural fields and pastures. The state of Mato Grosso do Sul is located inside the great South American open diagonal, a broad area of open formations extending from the northeastern Brazilian Caatinga to the southern Chaco in Argentina. The diagonal has a high herpetofauna biodiversity, including endemic species (Bucher 1980; Vanzolini 1988; Leynaud and Bucher 1999; Cabrera 1995, 1998; Duellman 1999; Colli *et al.* 2002; Souza 2005; Brusquetti and Lavilla 2006). The Chaco, one of the phytogeographic domains found in the diagonal, occurs in Mato Grosso do Sul in the southwestern municipality of Porto Murtinho.

The Chaco is a flat formation lying on top of the great Andean outwash plain, and encompasses approximately 1,000,000 km² (Bucher and Huszar 1999; Pennington *et al.* 2000). Chacoan phytophysionomies include shrubs and mesophytic vegetation usually associated with saline soil, and a great diversity of Leguminosae, Bromeliaceae, and Cactaceae (Pennington *et al.* 2000; Silva *et al.* 2000; Pott and Pott 2003). Climate is markedly seasonal, with a hot (over 40 °C) and wet summer and severe frosts in the dry winter (Pennington *et al.* 2000). The Chaco in Brazil (Porto Murtinho) is restricted to a small narrow strip located between the Paraguay and Apa rivers, not easily identified and defined in regional scale maps (Prado *et al.* 1992; Prado 1993). With the flat topography, the semi-arid climate, and the distinct plant formations (Prado *et al.* 1992; Prado 1993; Navarro *et al.* 2006), the Chaco is a well-defined biogeographical province (Morrone 2006) with associated endemism (Gallardo 1979; Leynaud and Bucher 1999; Pennington *et al.* 2000; Vinke and Vinke

2001; Morrone *et al.* 2004; Spichiger *et al.* 2004; Brusquetti and Lavilla 2006; Almeida *et al.* 2007).

At present, approximately 720 reptile species (Bérnils 2010) and 880 anuran species (SBH 2010) are known from Brazil. Most of the records on “Chacoan” herpetofauna are limited to Paraguay (Krieg 1948; Scott and Lovett 1975; Norman 1994; Ziegler *et al.* 2002) and Argentina (Cruz *et al.* 1992; Cabrera 1995, 1998; Cruz 1997) (readers must refer to Leynaud and Bucher 1999, Giraudo 2004 and Brusquetti and Lavilla 2006 for a broader literature review). For Brazil, the only known information about the Chacoan herpetofauna comes from the *Plano de Conservação da Bacia do Alto Paraguai* (PCBAP 1997), which cites 11 amphibian and four reptilian species, including three crocodilians. With this study we present a list of amphibians and reptiles from the Brazilian Chaco.

MATERIALS AND METHODS

Sampling was conducted bi-monthly from February 2008 to December 2009 throughout the municipality of Porto Murtinho, southwestern Mato Grosso do Sul state (approx. 21°40' S, 57°54' W to 21°56' S, 57°34' W; Figures 1 and 2). Sampling techniques included pitfall traps with drift fences, time-limited searches, and occasional encounters (Martins and Oliveira 1998). Pitfall trap consisted of 10 Y lines with four 60 L buckets (see Cechin and Martins 2000), each Y arm with 10 m long and each pitfall trap line distant 300 meters. Pitfall traps were kept opened for five days during sampling months. Voucher specimens were immediately killed with xylocaine, fixed in 5 % formalin, and transferred to 70 % alcohol.

The expected local species richness was estimated according to individual-based rarefaction curves for sampled localities, obtained in EstimateS 7.5 (Colwell 2005) with 1,000 randomizations of the original field samples employing the non-parametric Jackknife 1 index.

Specimens were deposited at the Coleção Zoológica




FIGURE 1. The major Paraguay ecoregions (modified from Cacciali 2010) neighboring Brazil and the close association of Porto Murtinho region with the Chaco biome (A). Points represent approximated herpetofauna sampling sites in Porto Murtinho Chaco region, state of Mato Grosso do Sul (B).




FIGURE 2. The dry (A) and wet (B) seasons create distinct physiognomies in Chaco throughout the year. (Photos: F.L. Souza).




FIGURE 3. Typical amphibian species from the region: *Leptodactylus bufonius* (A) and a juvenile *Ceratophrys cranwelli* (B). (Photos: L. Piatti).

de Referência of the Universidade Federal de Mato Grosso do Sul (ZUFMS, Campo Grande, Mato Grosso do Sul state). Collecting activities were carried out under permit IBAMA 10379.

RESULTS AND DISCUSSION

A total of 34 amphibian and 39 reptile species were registered in the Chaco of Porto Murtinho (Figures 3 and 4; Table 1). Hylidae (N = 13 species) was the most speciose amphibian family, and Dipsadidae was the richest reptilian family (N = 15 species).

Although rarefaction curves showed 36 amphibians (Figure 5A) and 54 reptile species could be expected (Figure 5B), the number of species recorded in the present study is similar to the projected herpetofauna richness for neighboring Cerrado (Diniz-Filho *et al.* 2007, 2009). Some typically Chacoan amphibian species, such as *Lepidobatrachus laevis* and *Leptodactylus laticeps*, recorded from neighboring Paraguayan Chaco areas (Ziegler *et al.* 2002; Brusquetti and Lavilla, 2006), were not found in Porto Murtinho. On the other hand, *Liophis poecilogyrus caesius*, a subspecies typical of the Chaco (Leynaud and Bucher 1999), was very common around Porto Murtinho and intergrades with *L. poecilogyrus schotti* from Cerrado in eastern Mato Grosso do Sul (Dixon and Marquezich 1992; Giraudo 2004). *Micrablepharus maximiliani*, a common

reptile species further east in the Brazilian Cerrado (Nogueira *et al.* 2009) was not found in Porto Murtinho. Both *L. poecilogyrus schotti* and *M. maximiliani* are found at Serra da Bodoquena, around 100 km east from Porto Murtinho (Uetanabaro *et al.* 2007; Nogueira *et al.* 2009). These results demonstrate the importance of Porto Murtinho as a real Chacoan province.

Most of the species can be considered open area dwellers reflecting partially the characteristics of this habitat. For amphibians, the flooding during the rainy season creates temporary ponds and favors their reproductive behavior. Also, some natural depressions accumulate permanent water and may represent important micro habitat during dry periods. Flooded areas are used by reptiles, such as *Acanthochelys macrocephala* (captured during rainy days), or they may favor the establishment of local aquatic populations of *Caiman yacare* and *Hydrodynastes gigas*. For these open area inhabitants, some species are typical of dry areas, such as *Liophis dilepis* (Caatinga), *Teius teyou/Leptodactylus bufonius* (Chaco), *Eupemphix nattereri* (Cerrado) (Brusquetti and Lavilla 2006; Ribeiro *et al.* 2008; Nogueira *et al.* 2009; Álvarez *et al.* 2009), showing that the Porto Murtinho Chaco has herpetological affinities with other open biomes. Finally, several species benefit from anthropic habitats and associated edge effects from the region (*Rhinella schneideri*, *Hypsiboas raniceps*, *Ameiva ameiva*), and have a broad geographical distribution in South America.



FIGURE 4. Typical reptile species from the region: *Liophis poecilogyrus caesius* (A) and *Teius teyou* (B). (Photos: F.L. Souza).




FIGURE 5. Rarefaction curves for amphibians (A) and reptiles (B) from Porto Murtinho Chaco region, Mato Grosso do Sul state. Curves represent mean richness (bars denote standard deviations) estimated by Jackknife 1 method after 1,000 randomizations of the sampled individuals.

TABLE 1. Checklist of amphibians and reptiles from Porto Murtinho Chaco region, state of Mato Grosso do Sul, recorded between February 2008 and December 2009.

SAMPLING SITES COORDINATES								
	22°04'54"	22°01'35"	22°04'49"	22°00'13"	21°54'29"	21°42'07"	21°39'00"	21°33'16"
	57°56'22"	57°54'15"	57°50'21"	57°46'26"	57°50'12"	57°43'13"	57°42'43"	57°46'51"
Bufonidae								
Melanophryniscus fulvoguttatus (Mertens, 1937)						X		X
Rhinella bergi (Céspedes, 2000 “1999”)	X	X	X	X	X	X	X	X
Rhinella major (Müller and Helmich, 1936)	X	X	X	X	X	X	X	X
Rhinella schneideri (Werner, 1894)	X	X	X			X		
Ceratophryidae								
Ceratophrys cranwelli Barrio, 1980						X		
Cycloramphidae								
Odontophrynus americanus (Duméril and Bibron, 1841)			X			X	X	
Hylidae								
Dendropsophus melanargyreus (Cope, 1887)						X		
Dendropsophus nanus (Boulenger, 1889)	X	X	X	X	X	X	X	X
Hypsiboas punctatus (Schneider, 1799)	X	X	X	X	X	X	X	X
Hypsiboas raniceps Cope, 1862	X	X	X	X	X	X	X	X
Phyllomedusa azurea Cope, 1862	X	X	X	X	X	X	X	X
Phyllomedusa sauvagii Boulenger, 1882		X	X			X	X	
Pseudis limellum (Cope, 1862)	X	X	X	X	X	X	X	X
Pseudis platensis Gallardo, 1961	X	X	X	X	X	X	X	X
Scinax acuminatus (Cope, 1862)	X	X	X	X	X	X	X	
Scinax fuscomarginatus (A. Lutz, 1925)						X	X	
Scinax fuscovarius (A. Lutz, 1925)	X	X	X			X		
Scinax nasicus (Cope, 1862)	X	X	X		X	X	X	X
Trachycephalus venulosus (Laurenti, 1768)		X	X			X		
Leiuperidae								
Eupemphix nattereri Steindachner, 1863						X		
Physalaemus albonotatus (Steindachner, 1864)	X	X	X	X	X	X	X	X
Physalaemus biligonigerus (Cope, 1861 “1860”)	X	X	X	X	X	X	X	X
Pseudopaludicola falcipes (Hensel, 1867)	X	X	X	X	X	X	X	X
Leptodactylidae								
Leptodactylus bufonius Boulenger, 1894	X	X	X	X	X	X	X	X
Leptodactylus chaquensis Cei, 1950	X	X	X	X	X	X	X	X
Leptodactylus diptyx Boettger, 1885		X	X		X	X		X
Leptodactylus elenae Heyer, 1978	X	X	X	X	X	X	X	X
Leptodactylus fuscus (Schneider, 1799)		X	X		X	X		
Leptodactylus labyrinthicus (Spix, 1824)						X		
Leptodactylus podicipinus (Cope, 1862)								
Microhylidae								
Chiasmocleis albopunctata (Boettger, 1885)				X				
Dermatonotus muelleri (Boettger, 1885)						X		X
Elachistocleis bicolor (Valenciennes in Guérin-Ménéville, 1838)		X	X		X	X	X	X
Elachistocleis ovalis (Schneider, 1799)		X	X			X	X	
Testudinidae								
Chelonoidis carbonaria (Spix, 1824)						X		
Chelidae								
Acanthochelys macrocephala Rhodin, Mittermeier and McMorris, 1984							X	
Phrynops geoffroanus (Schweigger, 1812)							X	
Alligatoridae								
Caiman yacare (Daudin, 1802)	X	X	X	X	X	X	X	X
Amphisbaenidae								
Amphisbaena cf. leseri			X					
Amphisbaena mertensii Strauch, 1881			X			X		
Iguanidae								
Iguana iguana (Linnaeus, 1758)	X							

TABLE 1. CONTINUED.

SAMPLING SITES COORDINATES								
	22°04'54"	22°01'35"	22°04'49"	22°00'13"	21°54'29"	21°42'07"	21°39'00"	21°33'16"
	57°56'22"	57°54'15"	57°50'21"	57°46'26"	57°50'12"	57°43'13"	57°42'43"	57°46'51"
Polychrotidae								
<i>Anolis meridionalis</i> Boettger, 1885						X		
<i>Anolis nitens</i> (Wagler, 1830)						X		
Tropiduridae								
<i>Tropidurus guarani</i> (Cope, 1862)			X			X		
Gekkonidae								
<i>Hemidactylus mabouia</i> (Moreau de Jonnès, 1818)	X		X			X		
Phyllodactylidae								
<i>Phyllopezus pollicaris</i> (Spix, 1825)			X			X		
Teiidae								
<i>Ameiva ameiva</i> (Linnaeus, 1758)	X	X	X	X	X	X	X	X
<i>Teius teyou</i> (Daudin, 1802)						X		X
<i>Tupinambis merianae</i> (Duméril and Bibron, 1839)	X	X	X		X	X		X
Gymnophthalmidae								
<i>Cercosaura ocellata</i> Wagler, 1830			X			X		
Scincidae								
<i>Mabuya frenata</i> (Cope, 1862)						X		X
<i>Mabuya guaporicola</i> Dunn, 1936						X		X
Boidae								
<i>Boa constrictor</i> Linnaeus, 1758						X		
<i>Eunectes notaeus</i> Cope, 1862	X					X		
Colubridae								
<i>Mastigodryas bifossatus</i> (Raddi, 1820)						X		
Dipsadidae								
<i>Helicops leopardinus</i> (Schlegel, 1837)						X		
<i>Hydrodynastes gigas</i> (Duméril, Bibron and Duméril, 1854)			X			X		X
<i>Leptodeira annulata</i> (Linnaeus, 1758)			X		X	X		X
<i>Liophis dilepis</i> (Cope, 1862)						X		
<i>Liophis poecilogyrus</i> (Wied, 1825)	X	X	X		X	X		X
<i>Liophis typhlus</i> (Linnaeus, 1758)						X		
<i>Mussurana bicolor</i> (Peracca, 1904)						X		
<i>Oxyrhopus rhombifer</i> Duméril, Bibron and Duméril, 1854						X		
<i>Philodryas patagoniensis</i> (Girard, 1858)			X			X		
<i>Pseudoboa nigra</i> (Duméril, Bibron and Duméril, 1854)						X		X
<i>Psomophis genimaculatus</i> (Boettger, 1885)						X		
<i>Thamnodynastes chaquensis</i> Bergna and Alvarez, 1993			X			X		X
<i>Thamnodynastes hypoconia</i> (Cope, 1860)						X		
<i>Xenodon matogrossensis</i> (Scrocchi and Cruz, 1993)						X		
<i>Xenodon merremii</i> (Wagler, 1824)						X		
Elapidae								
<i>Micrurus tricolor</i> Hoge, 1956						X		
Viperidae								
<i>Bothropoides mattogrossensis</i> (Amaral, 1925)			X		X	X	X	X
<i>Caudisona durissa</i> (Linnaeus, 1758)		X	X			X		

ACKNOWLEDGMENTS: The authors are grateful to Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq (Process 470484/2007-1) for financial support and to the owners who kindly permitted access to their farms, with special thanks to Oswaldo Ovelar (Seu Bala) and Fazenda Santa Vergínia staff. Norman J. Scott and C. Nogueira provided important comments on the manuscript. Thanks are extended to Otavio A. V. Marques, Ricardo J. Sawaya, C. Nogueira, G. Scrocchi, Daniel F. Silva, Jorge A. Céspedes, R. Montero, Paula H. Valdujo, and N. Jorge for help with species identification. F.L. Souza receives research grant from Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq (306034/2008-5).

LITERATURE CITED
Almeida, F.C., C.R. Bonvicino and P. Cordeiro-Estrela, P. 2007. Phylogeny and temporal diversification of *Calomys* (Rodentia, Sigmodontinae): implications for the biogeography of an endemic genus of the open/dry biomes of South America. *Molecular Phylogenetics and Evolution* 42(2): 449-466.
Álvarez, B.B., J.A.R. García, J.A. Céspedes, A.B. Hernando, V.H. Zaracho, C.C. Calamante and R.H. Aguirre. 2009. Herpetofauna, provinces of Chaco and Formosa, Chaco Oriental region, north-eastern Argentina. *Check List* 5(1): 74-82.
Bérnils, R.S. (org.). 2010. *Brazilian reptiles – List of species*. Electronic database accessible at <http://www.sbherpetologia.org.br/>. Sociedade Brasileira de Herpetologia. Captured on 16 June 2010.

- Brusquetti, F. and E.O. Lavilla. 2006. Lista comentada de los anfibios de Paraguay. Cuadernos de *Herpetología*, 20(2): 3-79.
- Bucher, E.H. 1980. Ecología de la fauna chaqueña. Una revisión. *Ecosur* 7(4): 111-221.
- Bucher, E.H. and P.C. Huszar. 1999. Sustainable management of the Gran Chaco of South America: Ecological promise and economic constraints. *Journal of Environmental Management* 57(2): 99-108.
- Cabrera, M.A. 1995. Comparative composition of turtle species in four natural regions of the Chacoan domain, South America. *Anales del Museo de Historia Natural de Valparaíso* 23(1): 41-52.
- Cabrera, M.A. 1998. *Las Tortugas Continentales da Sudamérica Austral*. Córdoba: BR Cópias., 119 p.
- Cacciali, P. 2010. Chromatic variation in populations of *Xenodon merremi* (Serpentes: Dipsadidae) in Paraguay. *Acta Herpetologica* 5(1): 107-112.
- Cechin, S.Z. and M. Martins. 2000. Eficiência de armadilhas de queda (pitfall traps) em amostragem de anfíbios e répteis no Brasil. *Revista Brasileira de Zoologia* 17(3): 729-740.
- Colli, G.R. 2005. As origens e a diversificação da herpetofauna do Cerrado; p. 247-264 In A. Scariot, J.C. Souza-Silva and J.M. Felfili (ed.). *Cerrado: Ecologia, Biodiversidade e Conservação*. Brasília: Ministério do Meio Ambiente.
- Colwell, R.K. 2005. *EstimateS: Statistic estimation of species richness and shared species from samples. Version 7.5*. Accessible at <http://viceroy.eeb.uconn.edu/estimates>. Captured on 17 August 2007.
- Cruz, F.B. 1997. Reproductive activity in *Tropidurus etheridgei* in the semiarid Chaco of Salta, Argentina. *Journal of Herpetology* 31(3): 444-450.
- Cruz, F.B., M.G. Perotti and L.A. Fitzgerald. 1992. Lista de anfibios y reptiles colectados en una localidad del Chaco Salteno. *Acta Zoologica Lilloana* 42(1): 101-107.
- Diniz-Filho, J.A.F., L.M. Bini, M.P. Pinto, T.F.L.V.B. Rangel, P.C. Carvalho, S.L. Vieira and R.P. Bastos. 2007. Conservation biogeography of anurans in Brazilian Cerrado. *Biodiversity and Conservation* 16(4): 997-1008.
- Diniz-Filho, J.A.F., L.M. Bini, G. Oliveira, B.S. Barreto, M.M.F.P. Silva, L.C. Terribile, T.F.L.V.B. Rangel, M.P. Pinto, N.P.R. Sousa, L.C.G. Vieira, A.S. Melo, P.M. Júnior, C.M. Vieira, D. Blamires, R.P. Bastos, P. Carvalho, L.G. Ferreira, M.P.C. Telles, F.M. Rodrigues, D. Melo e Silva, N.J. Silva Júnior and T.N. Soares. 2009. Macroecologia, biogeografia e áreas prioritárias para conservação no Cerrado. *Oecologia Brasiliensis* 13(3): 470-497.
- Dixon, J.R. and A.L. Markezich. 1992. Taxonomy and geographic variation of *Liophis poecilopyrus* (Wied) from South America (Serpentes: Colubridae). *The Texas Journal of Science* 44(2): 131-166.
- Duellman, W.E. 1999. *Patterns of distribution of amphibians: a global perspective*. Baltimore and London: Johns Hopkins University Press.
- Gallardo, J.M. 1979. Composicion, distribucion y origen de la herpetofauna chaqueña; p. 299-307 In W.E. Duellman (ed.). *The South American herpetofauna: its origin, evolution, and dispersal. Monograph of the Museum of Natural History, The University of Kansas* 7.
- Giraudo, A. 2004. *Serpientes de la Selva Paranaense y del Chaco Húmedo*. Buenos Aires: L.O.L.A (Literature of Latin América). 285 p.
- Krieg, H. 1948. *Zwischen Anden und Atlantik. Reisen eines Biologen in Südamerika*. Munich: Carl Hanser Verlag. 492 p.
- Leynaud, G.C., and E.H. Bucher. 1999. La fauna de serpientes del Chaco Sudamericano: diversidad, distribución geográfica y estado de conservación. *Academia Nacional de Ciencias de Córdoba. Miscelánea* 98: 1-45.
- Martins, M. and M.E. Oliveira. 1998. Natural history of snakes in forests of the Manaus region, central Amazonia, Brazil. *Herpetological Natural History* 6(2): 78-150.
- Morrone, J.J. 2006. Biogeographic areas and transition zones of latin america and the caribbean islands based on panbiogeographic and cladistic analyses of the entomofauna. *Annual Review of Entomology* 51: 467-494.
- Morrone, J.J., S.A. Mazzucconi and A.O. Bachmann. 2004. Distributional patterns of Chacoan water bugs (Heteroptera: Belostomatidae, Corixidae, Micronectidae and Gerridae). *Hydrobiologia* 523(1-3): 159-173.
- Navarro, G., J.A. Molina and L.P. Molas. 2006. Classification of the forests of the northern Paraguayan Chaco. *Phytocoenologia* 36(4): 473-508.
- Nogueira, C., G.R. Colli and M. Martins. 2009. Local richness and distribution of the lizard fauna in natural habitat mosaics of the Brazilian Cerrado. *Austral Ecology* 34(1): 83-96.
- Norman, D.R. 1994. *Anfibios y reptiles del Chaco Paraguayo –Tomo I. Amphibians and Reptiles of the Paraguayan Chaco – Volume I*. Heredia: Costa Rica. 281 p.
- PCBAP. 1997. Projeto Pantanal. Programa Nacional do Meio Ambiente (PNMA). *Subcomponente Pantanal. Plano de Conservação da Bacia do Alto Paraguai – PCBAP, v.2, t.3. Diagnóstico dos Meios físico e Biótico. Meio Biótico*. Brasília: MMA/PNMA. 433 p.
- Pennington, R.T., D.E. Prado and C.A. Pendry. 2000. Neotropical seasonally dry forests and Quaternary vegetation changes. *Journal of Biogeography* 27(2): 261-273.
- Pott, A. and V.J. Pott. 2003. Espécies de fragmentos florestais em Mato Grosso do Sul; p. 26-52 In R.B. Costa (org.). *Fragmentação Florestal e Alternativas de Desenvolvimento Rural na Região Centro-Oeste*. Campo Grande: Universidade Católica Don Bosco.
- Prado, D.E. 1993. What is the Gran Chaco vegetation in South America? I. A review. Contribution to the study of flora and vegetation of the Chaco, V. *Candollea* 48(1): 145-172.
- Prado, D.E., P.E. Gibbs, A. Pott and V.J. Pott. 1992. The Chaco-Pantanal transition in southern Mato Grosso, Brazil; p. 451-470. In P.A. Furley, J. Proctor and J.A. Ratter (ed.). *Nature and Dynamics of Forest-Savanna Boundaries*. London: Chapman and Hall.
- Ribeiro, S.C., F.S. Ferreira, S.V. Brito, G.G. Santana, W.L.S. Vieira, R.R.N. Alves, and W.O. Almeida. 2008. The squamata fauna of the Chapada do Araripe, Northeastern Brazil. *Cadernos de Cultura e Ciência* 1(1): 67-76.
- SBH. 2010. *Brazilian amphibians – List of species*. Electronic database accessible at <http://www.sbherpetologia.org.br>. Sociedade Brasileira de Herpetologia. Captured on 16 June 2010.
- Scott, N.J. and J.W. Lovett. 1975. A collection of reptiles and amphibians from the Chaco of Paraguay. *The University of Connecticut Occasional Papers, Biological Science Series* 2(16): 257-266.
- Silva Jr., N.J., C.A.D. Cintra, H.L.R. Silva, M.C. Costa, C.A. Souza, A.A. Pacheco Jr. and F.A. Gonçalves. 2009. Herpetofauna, Ponte de Pedra Hydroelectric Power Plant, states of Mato Grosso and Mato Grosso do Sul, Brazil. *Check List* 5(3): 518-525.
- Souza, F.L. 2005. Geographical distribution patterns of South American side-necked turtles (Chelidae), with emphasis on Brazilian species. *Revista Española de Herpetología* 19: 33-46.
- Spichiger, R., C. Calenge and B. Bise. 2004. Geographical zonation in the Neotropics of tree species characteristic of the Paraguay-Parana Basin. *Journal of Biogeography* 31(9): 1489-1501.
- Uetanabaro, M., F.L. Souza, P. Landgraf-Filho, A.F. Beda and R.A. Brandão. 2007. Anfíbios e répteis do Parque Nacional da Serra da Bodoquena, Mato Grosso do Sul, Brasil. *Biota Neotropica* 7(3): <http://www.biotaneotropica.org.br/v7n3/pt/abstract?inventory+bn01207032007>.
- Vanzolini, P.E. 1988. Distributional patterns of South American lizards; p. 317-342. In P.E. Vanzolini and W.R. Heyer (ed.). *Proceedings of a Workshop on Neotropical Distribution Patterns*. Rio de Janeiro: Academia Brasileira de Ciências.
- Vinke, T. and S. Vinke. 2001. The turtle and tortoise fauna of the central Chaco of Paraguay. *Radiata* 10(3): 3-19.
- Ziegler, T., J. Unger, A. Feiler and E. Lehr. 2002. The first Gran Chaco expedition of the Museum für Tierkunde Dresden: records of amphibians, reptiles, and mammals for the dry chaco of Paraguay (Amphibia, Reptilia, Mammalia). *Faunistische Abhandlungen, Staatliches Museum für Tierkunde Dresden* 23(11): 219-237.

RECEIVED: February 2010

REVISED: July 2010

ACCEPTED: August 2010

PUBLISHED ONLINE: September 2010

EDITORIAL RESPONSIBILITY: Alejandro R. Giraudo